



# Mark Scheme (Results)

January 2022

Pearson Edexcel International GCSE

In Physics (4PH1) Paper 1P and Science

(Double Award) (4SD0) Paper 1P

## **Edexcel and BTEC Qualifications**

Edexcel and BTEC qualifications are awarded by Pearson, the UK's largest awarding body. We provide a wide range of qualifications including academic, vocational, occupational and specific programmes for employers. For further information visit our qualifications websites at [www.edexcel.com](http://www.edexcel.com) or [www.btec.co.uk](http://www.btec.co.uk). Alternatively, you can get in touch with us using the details on our contact us page at [www.edexcel.com/contactus](http://www.edexcel.com/contactus).

## **Pearson: helping people progress, everywhere**

Pearson aspires to be the world's leading learning company. Our aim is to help everyone progress in their lives through education. We believe in every kind of learning, for all kinds of people, wherever they are in the world. We've been involved in education for over 150 years, and by working across 70 countries, in 100 languages, we have built an international reputation for our commitment to high standards and raising achievement through innovation in education. Find out more about how we can help you and your students at: [www.pearson.com/uk](http://www.pearson.com/uk)

January 2022

Question Paper Log Number P70705A

Publications Code 4PH1\_1P\_2201\_MS

All the material in this publication is copyright

© Pearson Education Ltd 2022

## General Marking Guidance

- All candidates must receive the same treatment. Examiners must mark the first candidate in exactly the same way as they mark the last.
- Mark schemes should be applied positively. Candidates must be rewarded for what they have shown they can do rather than penalised for omissions.
- Examiners should mark according to the mark scheme not according to their perception of where the grade boundaries may lie.
- There is no ceiling on achievement. All marks on the mark scheme should be used appropriately.
- All the marks on the mark scheme are designed to be awarded. Examiners should always award full marks if deserved, i.e. if the answer matches the mark scheme. Examiners should also be prepared to award zero marks if the candidate's response is not worthy of credit according to the mark scheme.
- Where some judgement is required, mark schemes will provide the principles by which marks will be awarded and exemplification may be limited.
- When examiners are in doubt regarding the application of the mark scheme to a candidate's response, the team leader must be consulted.
- Crossed out work should be marked UNLESS the candidate has replaced it with an alternative response.

Question number	Answer	Notes	Marks
1 (a)	B;  A is incorrect because the top two field lines are in the wrong direction C is incorrect because all the field lines are in the wrong direction D is incorrect because the second field line from the top is in the wrong direction		1
(b) (i)	B;  A is incorrect because there is a magnetic force on each of the magnets C is incorrect because the two nearest poles are opposite not alike D is incorrect because the forces on the magnets pass through their centres of gravity		1
(ii)	C;  A is incorrect because the field lines are not straight nor parallel to each other B is incorrect because the field lines are not straight nor parallel to each other D is incorrect because the field lines are not parallel to each other		1
(c)	steel / nickel / cobalt / neodymium;	condone iron	1
(d)	EITHER:  (plotting) compass used; multiple compasses or repeated use of single compass; joining up of dots/idea of compasses forming continuous line;  OR  iron filings used; sprinkled / eq;  card tapped (to reveal pattern);	all marks can be awarded from a diagram       ignore 'poured' or other heavy-handed method	3

Total for Question 1 = 7 marks

Question number	Answer	Notes	Marks
2 (a)	idea of subtracting the background count rate;		1
(b) (i)	time taken;  and either of for (radio)activity to halve; for half of the (radioactive) nuclei / atoms / isotope to decay;	allow "how long it takes" reject "half the time"  allow count rate for activity ignore mass, substance	2
(ii)	indication on graph of a half in count rate;  2.6 (minutes);	e.g. line drawn across from 25 until it reaches the curve, then down to the time axis allow 2.5-2.7 (minutes) 2.3 (minutes) = 1 mark	2

Total for Question 2 = 5 marks

Question number	Answer	Notes	Marks
3 (a)	(i) travel at the same speed (in a vacuum) / can travel in a vacuum;	allow both transverse waves, both transfer energy, both microwaves	1
	(ii) wavelength;	allow idea of different range / penetration ignore amplitude	1
(b)	(i) speed = frequency $\times$ wavelength;	allow standard symbols and rearrangements e.g. $\lambda = v / f$ allow c for speed ignore s for speed	1
	(ii) substitution OR rearrangement; evaluation;  e.g. $3.0 \times 10^8 = 5.2 \times 10^9 \times \lambda$ OR $\lambda = v / f$ (wavelength =) 0.058 (m)	-1 for POT error  allow 0.06, 0.0576...(m)	2
(c)	(i) D;  A is incorrect because electromagnetic waves are not longitudinal B is incorrect because electromagnetic waves are not mechanical waves C is incorrect because sound waves are not electromagnetic waves		1
	(ii) vibrations / oscillations; correct relationship between direction of travel/energy transfer and direction of vibration for both transverse and longitudinal waves;	both marks can be scored from a suitable diagram  allow movement of particles / eq for vibrations for this mark	2

Total for Question 3 = 8 marks

Question number	Answer	Notes	Marks
4 (a)	protractor;		1
(b) (i)	any indication that the angle is between the normal and the incident ray;		1
(ii)	79 (degrees);	allow 78-80 (degrees) ecf from indicated angle of incidence in (i) e.g. 10-12 (degrees) if angle marked between ray and boundary	1
(iii)	any straight ray to the right of the normal that comes from the point of incidence; correct angle of reflection;	judge by eye  judge by eye	2

Total for Question 4 = 5 marks

Question number	Answer	Notes	Marks
5 (a)	(i) (average) speed = distance (travelled) ÷ time (taken);	allow standard symbols and rearrangements e.g. $v = s / t$ allow s, d for distance condone s for speed	1
	(ii) substitution OR rearrangement; evaluation;  e.g. $21 = \text{distance} / 0.14$ OR $s = v \times t$ (distance =) 2.9 (m)	allow 3, 2.94 (m)	2
(b)	(i) force = mass × acceleration;	allow standard symbols and rearrangements e.g. $a = F / m$	1
	(ii) substitution OR rearrangement; evaluation;  e.g. $7600 = 1200 \times a$ OR $a = F / m$ (a =) (-)6.3 (m/s <sup>2</sup> )	allow 6.33... (m/s <sup>2</sup> )	2
	(iii) substitution into $v^2 = u^2 + 2as$ ; rearrangement; evaluation;  e.g. $0^2 = 21^2 + [2 \times (-)6.3 \times \text{distance}]$ distance = 441 / 12.6 distance = 35 (m)	ecf answer from (ii)  allow 34.8...(m)	3

Total for Question 5 = 9 marks



Question number	Answer	Notes	Marks
6 (a)	(i) any attempt to find gradient of graph;  use of two points on the line to calculate gradient;  evaluation;  e.g. acceleration = gradient acceleration = $(-)/4.2 / 0.45$ (acceleration =) $-9.3 \text{ (m/s}^2\text{)}$	allow use of acceleration formula allow reading of pair of velocities with matching time interval reject positive answer  allow $-9.3$ to $-9.4$	3
	(ii) any clear indication that distance travelled = area;  correct use of data from graph; evaluation;  e.g. distance = area distance = $0.5 \times 0.45 \times 4.2$ (distance =) $0.95 \text{ (m)}$	accept alternative method using $v^2 = u^2 + 2as$ with acceleration calculated in (i) allow attempt to calculate area of triangle  allow $0.94, 0.945 \text{ (m)}$	3
(b)	(i) weight / gravitational force;  drag / air resistance;	ignore unqualified 'gravity', gravitational field strength ignore upthrust, lift	2
	(ii) one upward arrow and one downward arrow drawn; arrows originate at object; downward arrow drawn longer than upward arrow;	judge by eye  judge by eye	3
	(iii) any four from: MP1. object is accelerating (from A to B); MP2. downward force greater than upward force (at A); MP3. gradient / acceleration decreasing (from A to B); MP4. drag increases as speed increases; MP5. resultant force decreases; MP6. idea that (just after) B, downward force = upward force; MP7. idea that in region BC, acceleration is zero/close to zero; MP8. terminal velocity achieved in region BC;	allow speeding up allow any recognisable upward force and downward force  allow any recognisable upward force and downward force  allow constant velocity	4

Total for Question 6 = 15 marks

Question number	Answer	Notes	Marks
7 (a) (i)	idea that voltage across thermistor + voltage across fixed resistor = voltage across cell; 0.59 (V);	allow 0.632 (V)	2
(ii)	voltage = current × resistance;	allow standard symbols and rearrangements e.g. V, I and R ignore c,C for current	1
(iii)	substitution; rearrangement; evaluation;  e.g. $0.59 = 0.0062 \times R$ $R = 0.59 / 0.0062$ (R =) 95 (Ω)	ecf answer from (i)  -1 for POT error  answers of R = 90.7... or R = 101.9...(Ω) gain full marks answer of 242 (Ω) gains 2 marks  allow 95.2, 95.16... condone 95.1	3
(b) (i)	idea that resistance of thermistor decreases with an increase in temperature; idea of non-linear relationship;	allow idea that rate of change is decreasing resistance inversely proportional to temperature scores both marks	2
(ii)	voltmeter reading decreases;  (because) resistance of <b>thermistor</b> increases; idea that current in circuit/thermistor decreases;	allow voltage across resistor decreases	3

Total for Question 7 = 11 marks

Question number	Answer	Notes	Marks														
8 (a)	(i) idea that kinetic store increases;	e.g. chemical transferred to kinetic	1														
	(ii) idea that gravitational store increases;		1														
(b)	two correct statements ticked;; <table border="1" data-bbox="358 495 842 814"> <thead> <tr> <th>Statement</th> <th>Correct (✓)</th> </tr> </thead> <tbody> <tr> <td>gravitational store increases</td> <td>✓</td> </tr> <tr> <td>gravitational store stays the same</td> <td></td> </tr> <tr> <td>gravitational store decreases</td> <td></td> </tr> <tr> <td>kinetic store increases</td> <td></td> </tr> <tr> <td>kinetic store stays the same</td> <td></td> </tr> <tr> <td>kinetic store decreases</td> <td>✓</td> </tr> </tbody> </table>	Statement	Correct (✓)	gravitational store increases	✓	gravitational store stays the same		gravitational store decreases		kinetic store increases		kinetic store stays the same		kinetic store decreases	✓	3 ticks scores 1 max 4 or more ticks scores 0	2
Statement	Correct (✓)																
gravitational store increases	✓																
gravitational store stays the same																	
gravitational store decreases																	
kinetic store increases																	
kinetic store stays the same																	
kinetic store decreases	✓																
(c)	(i) gravitational (force);	allow weight, gravity	1														
	(ii) substitution into given formula; evaluation;  e.g. orbital speed = $(2 \times \pi \times 7100) / 5800$ (orbital speed =) 7.7 (km/s)	allow 7.69... (km/s)	2														
(d)	any four from:  MP1. bars increase in temperature when facing towards Sun / decrease in temperature when facing away from Sun; MP2. (when pointed at the Sun,) black bar increases temperature faster than white bar; MP3. (because) black is a better absorber of radiation than white; MP4. (so) black bar reaches a higher temperature than white bar; MP5. (when pointed away from the Sun,) black bar decreases temperature faster than white bar; MP6. (because) black is a better emitter of radiation than white; MP7. convection/conduction plays no part in heat transfer (outside the spacecraft); MP8. (because) there are no particles outside the spacecraft;	accept any clear reverse argument  allow energy, heat, IR for radiation  allow energy, heat, IR for radiation	4														

Total for Question 8 = 11 marks

Question number	Answer	Notes	Marks																		
9 (a)	<p>any five from:</p> <p>MP1. measure original length of spring;  MP2. measure new length / extension for a <b>range</b> of masses;  MP3. extension = new length - original length;  MP4. use of ruler;  MP5. <b>method</b> of avoiding parallax, e.g. look at eye level or use a pointer;  MP6. use of a set square / clamping ruler vertically;  MP7. idea of measuring between the same two points (on the spring);  MP8. idea of repeating and averaging;  MP9. idea of measuring extension with decreasing mass as well;</p>	<p>allow any marking points if seen on diagram</p> <p>allow tape measure</p> <p>allow repeating to identify anomalies</p>	5																		
(b) (i)	<p>suitable linear scale chosen (&gt;50% of grid used); axes labelled with quantities and units;</p> <p>all plotting correct to nearest half square;</p>	<p>ignore orientation</p> <table border="1"> <thead> <tr> <th>Force in N</th> <th>Extension in cm</th> </tr> </thead> <tbody> <tr><td>0.0</td><td>0.0</td></tr> <tr><td>1.0</td><td>2.5</td></tr> <tr><td>2.0</td><td>5.0</td></tr> <tr><td>3.0</td><td>9.8</td></tr> <tr><td>4.0</td><td>10.0</td></tr> <tr><td>5.0</td><td>12.5</td></tr> <tr><td>6.0</td><td>15.5</td></tr> <tr><td>7.0</td><td>19.5</td></tr> </tbody> </table>	Force in N	Extension in cm	0.0	0.0	1.0	2.5	2.0	5.0	3.0	9.8	4.0	10.0	5.0	12.5	6.0	15.5	7.0	19.5	3
Force in N	Extension in cm																				
0.0	0.0																				
1.0	2.5																				
2.0	5.0																				
3.0	9.8																				
4.0	10.0																				
5.0	12.5																				
6.0	15.5																				
7.0	19.5																				
(ii)	data point at (3.0,9.8) identified;	allow (7.0,19.5) as chosen point	1																		
(iii)	straight line of best fit passing through origin and non-anomalous points;	ecf from plotting in (i) ignore line beyond 6.0N	1																		
(iv)	<p>any three from:</p> <p>MP1. quotation of Hooke's Law;  MP2. line is straight;  MP3. line passes through origin;  MP4. no evidence of having passed elastic limit;</p>	<p>i.e. force and extension should be proportional  allow idea that line is not straight if consistent with (iii)</p> <p>allow idea that elastic limit has been reached if consistent with (iii)</p>	3																		

Total for Question 9 = 13 marks

Question number	Answer	Notes	Marks
10 (a) (i)	36 (degrees);		1
(ii)	refractive index = $\sin(i) / \sin(r)$ ;	allow standard symbols and rearrangements e.g. $n = \sin(i) / \sin(r)$	1
(iii)	substitution; evaluation; answer quoted to 2 s.f.;	allow ecf from (i)  mark independently	3
	e.g. refractive index = $\sin(61) / \sin(36)$ (refractive index =) 1.48799... (refractive index =) 1.5		
(b)	red refracts less than violet;  correct link made between colour and refractive index;  correct link made between wavelength and refractive index;	allow RA allow red bends less than violet e.g. red has a lower refractive index than violet e.g. refractive index decreases with increasing wavelength	3

Total for Question 10 = 8 marks

Question number	Answer	Notes	Marks
11 (a) (i)	rearrangement OR substitution into given formula; evaluation;  e.g. $V_2 = p_1 \times V_1 / p_2$ OR $120 \times 92 = 64 \times V_2$ (volume =) 170 (m <sup>3</sup> )	allow 172, 173, 172.5	2
(ii)	constant temperature / amount of air / mass of air;	however expressed e.g. number of particles constant	1
(b) (i)	any three from: MP1. (reduction in temperature) reduces speed/KE of particles; MP2. idea of fewer collisions with walls per unit time; MP3. idea of each collision with wall being less 'hard'; MP4. force (per unit area) on the container decreases;	allow particles collide with walls less often	3
(ii)	substitution into given formula; rearrangement; evaluation;  e.g. $120 / 290 = 64 / T_2$ $T_2 = (64 \times 290) / 120$ (temperature =) 150 (K)	allow 155, 154.6... (K)	3

Total for Question 11 = 9 marks

Question number	Answer	Notes	Marks
12 (a)	<p>calculation of energy transferred by battery; efficiency formula stated;</p> <p>correct substitution;</p> <p>evaluation;</p> <p>e.g. energy supplied = <math>VIt = 12 \times 0.25 \times 12 = 36</math> (J) efficiency = <math>\frac{\text{useful energy output}}{\text{total energy output}}</math> efficiency = <math>25 / 36</math> (<math>\times 100\%</math>) efficiency = 69 (%)</p>	<p>36 (J) seen seen or implied anywhere in working allow ecf from battery energy if clear <math>25/36</math> (<math>\times 100</math>) seen</p> <p>allow 70, 69.4...(%)</p>	4
(b) (i)	<p>current / coil has a magnetic field; interaction between fields; resulting in a force;</p> <p>forces on opposite sides of the coil are in opposite directions;</p>	ignore references to attraction / repulsion	4
(ii)	<p>C - YZ;</p> <p>A is incorrect because WX moves downwards B is incorrect because part of XY moves downwards D is incorrect because part of ZW moves downwards</p>		1

Total for Question 12 = 9 marks

