

Please check the examination details below before entering your candidate information



Candidate surname

Other names

**Pearson Edexcel**  
International  
Advanced Level

Centre Number

--	--	--	--	--

Candidate Number

--	--	--	--	--

**Friday 11 October 2019**

Morning (Time: 1 hour 30 minutes)

Paper Reference **WCH12/01**

**Chemistry**

**International Advanced Subsidiary Level**

**Unit 2: Energetics, Group Chemistry, Halogenoalkanes and Alcohols**

**Candidates must have: Scientific calculator  
Data Booklet  
Ruler**

Total Marks

### Instructions

- Use **black** ink or **black** ball-point pen.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions.
- Answer the questions in the spaces provided – *there may be more space than you need.*

### Information

- The total mark for this paper is 80.
- The marks for **each** question are shown in brackets – *use this as a guide as to how much time to spend on each question.*
- In the question marked with an **asterisk** (\*), marks will be awarded for your ability to structure your answer logically, showing how the points that you make are related or follow on from each other where appropriate.
- There is a Periodic Table on the back cover of this paper.

### Advice

- Read each question carefully before you start to answer it.
- Show all your working in calculations and include units where appropriate.
- Check your answers if you have time at the end.

Turn over ►

P61657A

©2019 Pearson Education Ltd.

1/1/1/1/1/1/



Pearson


## SECTION A

Answer ALL the questions in this section.

You should aim to spend no more than 20 minutes on this section.

For each question, select one answer from A to D and put a cross in the box .  
If you change your mind, put a line through the box  and then mark your new answer with a cross .

- 1 Lithium carbonate decomposes on heating.



What is the maximum volume, in  $\text{dm}^3$ , measured at room temperature and pressure (r.t.p.), of gas produced from 3.69 g of lithium carbonate?

$[M_r \text{Li}_2\text{CO}_3 = 73.8 \quad \text{Molar volume of a gas at r.t.p.} = 24.0 \text{ dm}^3 \text{ mol}^{-1}]$

- A 24.0  
 B 12.0  
 C 1.20  
 D 0.83

(Total for Question 1 = 1 mark)

- 2 A sample of 1,2-dichloroethane,  $\text{CH}_2\text{ClCH}_2\text{Cl}$ , contains only the isotopes  $^1\text{H}$ ,  $^{12}\text{C}$ ,  $^{35}\text{Cl}$  and  $^{37}\text{Cl}$ .

How many molecular ion peaks are there in its mass spectrum?

- A 1  
 B 2  
 C 3  
 D 4

(Total for Question 2 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



3 Under certain conditions, graphite burns to form carbon monoxide.



Which of these is correct?

- A  $\Delta_c H$  (carbon) =  $-221 \text{ kJ mol}^{-1}$
- B  $\Delta_f H$  (carbon monoxide) =  $-221 \text{ kJ mol}^{-1}$
- C  $\Delta_c H$  (carbon) =  $-110.5 \text{ kJ mol}^{-1}$
- D  $\Delta_f H$  (carbon monoxide) =  $-110.5 \text{ kJ mol}^{-1}$

(Total for Question 3 = 1 mark)

4 What are the strongest interactions **between** molecules in solid hydrogen iodide, HI?




- A covalent bonds
- B hydrogen bonds
- C ionic bonds
- D London forces

(Total for Question 4 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



5 Equations for four reactions of copper or its compounds are shown.



(a) Which is a disproportionation reaction?

(1)

- A Reaction 1
- B Reaction 2
- C Reaction 3
- D Reaction 4

(b) Which is an acid-base reaction?

(1)

- A Reaction 1
- B Reaction 2
- C Reaction 3
- D Reaction 4

(Total for Question 5 = 2 marks)

6 Which statement is correct?

- A barium carbonate is less stable to heat than magnesium carbonate
- B barium hydroxide is less soluble in water than magnesium hydroxide
- C barium sulfate is less soluble in water than magnesium sulfate
- D barium metal is less reactive with water than magnesium metal

(Total for Question 6 = 1 mark)

Use this space for any rough working. Anything you write in this space will gain no credit.



7 Which statement is **not** correct?

- A chlorine is more electronegative than bromine
- B chlorine is more reactive than bromine
- C chloride ions are stronger reducing agents than bromide ions
- D chloride ions are stronger reducing agents than fluoride ions

(Total for Question 7 = 1 mark)

8 A white solid **X** produces a red colour in a flame test.

When aqueous silver nitrate and nitric acid are added to a solution of **X**, a cream precipitate is formed which dissolves in concentrated aqueous ammonia.

What is the formula of **X**?

- A  $\text{SrBr}_2$
- B  $\text{NaBr}$
- C  $\text{LiCl}$
- D  $\text{BaI}_2$

(Total for Question 8 = 1 mark)

9 Exactly  $50.0 \text{ cm}^3$  of  $2.00 \text{ mol dm}^{-3}$  nitric acid reacts with  $50.0 \text{ cm}^3$  of  $1.00 \text{ mol dm}^{-3}$  barium hydroxide to form a neutral solution of barium nitrate.

(a) What is the concentration, in  $\text{mol dm}^{-3}$ , of barium nitrate in the solution?

(1)

- A 0.05
- B 0.50
- C 1.00
- D 2.00

(b) The volume of the nitric acid is measured using a burette. Each burette reading has an uncertainty of  $\pm 0.05 \text{ cm}^3$ .

What is the percentage uncertainty in measuring  $50.00 \text{ cm}^3$  of the nitric acid?

(1)

- A  $\pm 0.40\%$
- B  $\pm 0.20\%$
- C  $\pm 0.10\%$
- D  $\pm 0.05\%$

(Total for Question 9 = 2 marks)





13 The equation for a reversible reaction is shown.



What effect will each change have on the rate of reaction and the equilibrium yield of phosphorus(V) chloride?

(a) Increasing the temperature at constant pressure.

(1)

	Effect on rate of reaction	Effect on yield of $\text{PCl}_5(\text{g})$
<input type="checkbox"/> A	increase	decrease
<input type="checkbox"/> B	decrease	decrease
<input type="checkbox"/> C	increase	increase
<input type="checkbox"/> D	decrease	increase

(b) Increasing the pressure at constant temperature.

(1)

	Effect on rate of reaction	Effect on yield of $\text{PCl}_5(\text{g})$
<input type="checkbox"/> A	increase	decrease
<input type="checkbox"/> B	decrease	decrease
<input type="checkbox"/> C	increase	increase
<input type="checkbox"/> D	decrease	increase

(Total for Question 13 = 2 marks)

14 A chloroalkane is heated with dilute aqueous sodium hydroxide and the pure organic product is obtained.

When the organic product is warmed with acidified potassium dichromate(VI) solution, there is no change in colour.

The chloroalkane could be

- A 1-chlorobutane
- B 2-chloro-2-methylpropane
- C 1-chloro-2-methylpropane
- D 2-chlorobutane

(Total for Question 14 = 1 mark)



15 This question is about two isomeric alcohols and two isomeric carbonyl compounds.

propan-1-ol,  $\text{CH}_3\text{CH}_2\text{CH}_2\text{OH}$  and propan-2-ol,  $\text{CH}_3\text{CH}(\text{OH})\text{CH}_3$

propanal,  $\text{CH}_3\text{CH}_2\text{CHO}$  and propanone,  $\text{CH}_3\text{COCH}_3$

(a) Which reaction is possible?

(1)

- A reducing propan-1-ol to propanal
- B oxidising propan-1-ol to propanal
- C reducing propanal to propanone
- D oxidising propan-1-ol to propanone

(b) Which compound would be expected to give a significant peak at  $m/z = 31$  in its mass spectrum?

(1)

- A propan-1-ol
- B propan-2-ol
- C propanal
- D propanone

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





(c) The infrared spectrum of one of the four compounds is shown.



Identify the compound, using the infrared absorptions from the Data Booklet.

(1)

- A propan-1-ol
- B propan-2-ol
- C propanal
- D propanone

(Total for Question 15 = 3 marks)

**TOTAL FOR SECTION A = 20 MARKS**





## SECTION B

Answer ALL the questions. Write your answers in the spaces provided.

16 Group 2 hydroxides,  $M(OH)_2$ , are used to neutralise acids.

- (a) Write an equation for the reaction of calcium with cold water.  
State symbols are not required.

(1)

- (b) Explain why a saturated solution of calcium hydroxide is more alkaline than a saturated solution of magnesium hydroxide.

(2)

.....

.....

.....

.....

.....

.....

- (c) A solution was tested for the presence of carbonate ions.

Dilute hydrochloric acid was added to the solution and the gas given off was bubbled through limewater (aqueous calcium hydroxide). A white precipitate formed.

- (i) Write the **ionic** equation for the formation of the gas.  
State symbols are not required.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA



- (ii) Write the equation for the formation of the precipitate.  
Include state symbols.

(2)

- (d) Magnesium hydroxide can be used to neutralise hydrochloric acid in the stomach to relieve acid indigestion.



Calculate the minimum mass of magnesium hydroxide needed to neutralise 0.150 mol of hydrochloric acid.

Give your answer to an appropriate number of significant figures.

(3)




(Total for Question 16 = 9 marks)



17 Biobutanol is a possible alternative to bioethanol as an additive to petrol.

(a) (i) Complete the table for the four isomers of  $C_4H_9OH$  which are alcohols.

(3)

Skeletal formula	Name	Classification	Boiling temperature / °C
	butan-1-ol	primary	117
			108
			99
	2-methylpropan-2-ol		82

(ii) Explain the difference in boiling temperature between butan-1-ol and 2-methylpropan-2-ol.

(2)

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(iii) The displayed formula of butan-1-ol is shown.

Complete the diagram to show the strongest intermolecular force between **two** molecules of butan-1-ol. Include the intermolecular bond angle.

(2)



DO NOT WRITE IN THIS AREA


DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) (i) Complete the equation for the combustion of butan-1-ol.  
State symbols are not required.

(1)



(ii) Calculate a value for the enthalpy change of combustion of butan-1-ol using the equation in (b)(i) and the mean bond enthalpies in the table.

(3)

Bond	C—C	C—O	C=O	C—H	O—H	O=O
Mean bond enthalpy / kJ mol <sup>-1</sup>	347	358	805	413	464	498

(iii) A data book value of the molar enthalpy change of combustion of butan-1-ol is  $-2670 \text{ kJ mol}^{-1}$

Give **two** reasons for the difference between this value and the value calculated in (b)(ii).

(2)

.....

.....

.....

.....

.....

(c) Biobutanol has some advantages over bioethanol.

(i) The combustion of bioethanol releases  $23 \text{ MJ dm}^{-3}$  compared to petrol which releases  $32 \text{ MJ dm}^{-3}$ .

Calculate the energy released in  $\text{MJ dm}^{-3}$  for the combustion of biobutanol.

Biofuel	Formula	$\Delta_c H / \text{kJ mol}^{-1}$	Density / $\text{g cm}^{-3}$
biobutanol	$\text{C}_4\text{H}_9\text{OH}$	$-2670$	$0.810$

(2)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





(ii) Biobutanol can be mixed with petrol in any proportion whereas bioethanol cannot.

Petrol is a mixture of liquid alkanes.

Explain why petrol is more miscible with biobutanol than with bioethanol.

(2)

.....

.....

.....

.....

.....

.....

.....

**(Total for Question 17 = 17 marks)**

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA







DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA


**BLANK PAGE**




18 Two different processes can be used for the production of chloroethene ( $\text{CH}_2=\text{CHCl}$ ), which is the monomer for the manufacture of poly(chloroethene).

(a) Process A

Ethene, produced from crude oil, reacts with chlorine gas to form 1,2-dichloroethane. The 1,2-dichloroethane is then thermally cracked to form chloroethene.



(i) A student wrote a mechanism for Reaction 1.



The mechanism contains two mistakes.

Identify each mistake, giving the correction that should be made.

(2)

Mistake 1.....

.....

Correction.....

.....

Mistake 2.....

.....

Correction.....

.....



(ii) Write the overall equation for Process **A** and use it to calculate the percentage atom economy by mass for the production of chloroethene.

(3)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(b) Process **B**

In an alternative process, chloroethene is obtained by reacting ethyne ( $\text{HC}\equiv\text{CH}$ ), with hydrogen chloride. Ethyne is produced from coal. This reaction uses a catalyst of mercury(II) chloride which is highly toxic.



The temperature during this reaction is high enough to vaporise the mercury(II) chloride catalyst.

State how Process **A** compares to Process **B** in terms of the percentage atom economy by mass and environmental impact of each process. No calculation is needed.

(2)

.....

.....

.....

.....

.....

.....

.....

.....





\*(c) Process **B** involves gas molecules reacting at high temperature in the presence of a catalyst.

Explain the effects on the rate of reaction of increasing the temperature and using a catalyst in Process **B**, referring to a labelled diagram of the Maxwell-Boltzmann distribution.

(6)

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA





DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

Handwriting practice area with horizontal dotted lines.

**(Total for Question 18 = 13 marks)**

**TOTAL FOR SECTION B = 39 MARKS**



P 6 1 6 5 7 A 0 2 1 2 8



DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

**BLANK PAGE**




## SECTION C

Answer ALL the questions. Write your answers in the spaces provided.

19 Major volcanic eruptions, such as the one in Indonesia in 1815, eject large amounts of ash and gases into the atmosphere. The gases include carbon dioxide, water vapour, hydrogen chloride and sulfur dioxide.

- (a) (i) Complete the dot-and-cross diagram to show a possible arrangement of the outer shell electrons in a molecule of sulfur dioxide.  
Use dots (•) for the sulfur electrons and crosses (×) for the oxygen electrons.

(2)



- (ii) Suggest a value for the bond angle.

(1)

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



(b) The levels of sulfur dioxide in the atmosphere can be measured by collection and titration with sodium hydroxide solution.

One possible method is:

- air containing sulfur dioxide is bubbled through hydrogen peroxide solution
- all the sulfur dioxide reacts to form 40.0 cm<sup>3</sup> of dilute sulfuric acid, solution Z



- a pipette is used to remove 10.0 cm<sup>3</sup> portions of solution Z
- each portion is titrated with 0.00500 mol dm<sup>-3</sup> sodium hydroxide.

The results are shown in the table.


Titration	1	2	3
Final volume/cm <sup>3</sup>	21.55	42.70	21.05
Initial volume/cm <sup>3</sup>	0.00	21.55	0.00
Titre/cm <sup>3</sup>	21.55	21.15	21.05

The mean titre is 21.10 cm<sup>3</sup>

(i) Give a reason why a further titration was not attempted.

(1)

(ii) Calculate the number of moles of sulfuric acid in the 40 cm<sup>3</sup> of solution Z.




(3)





- (iii) Deduce the number of moles of sulfur dioxide bubbled through the hydrogen peroxide solution, using the answer from (b)(ii) and the equation





- (iv) The air containing sulfur dioxide was bubbled through the hydrogen peroxide solution at a rate of  $10 \text{ dm}^3 \text{ min}^{-1}$  for 30 minutes.

Calculate the concentration of sulfur dioxide in the air in parts per million (ppm) by volume.

The molar volume of a gas at r.t.p. is  $24 \text{ dm}^3 \text{ mol}^{-1}$ . (3)

- (c) (i) During a volcanic eruption, hydrogen chloride gas is also released into the upper atmosphere, which in turn produces some chlorine free radicals. Chlorine free radicals react with ozone:



Derive the overall equation for this reaction of ozone.  
State symbols are not required.

(1)



(ii) Give **two** reasons why the presence of a small number of chlorine free radicals in the upper atmosphere causes a large decrease in the amount of ozone.

(2)

.....

.....

.....


.....

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

(d) (i) Sulfur dioxide is converted into sulfur trioxide in the upper atmosphere in the presence of ultraviolet light.



Show, by use of all the relevant oxidation numbers, that this is a redox reaction.

(2)


.....

.....

.....


.....

(ii) Sulfur trioxide reacts with water to produce sulfuric acid.



Draw an enthalpy level diagram to show the enthalpy change for this reaction.

(2)





# The Periodic Table of Elements

1 2 3 4 5 6 7 0 (8) (18)

1.0	<b>H</b>
	hydrogen
	1

## Key

relative atomic mass
<b>atomic symbol</b>
name
atomic (proton) number

(1) (2)

6.9	<b>Li</b>	lithium	3
9.0	<b>Be</b>	beryllium	4
23.0	<b>Na</b>	sodium	11
24.3	<b>Mg</b>	magnesium	12

(13)

10.8	<b>B</b>	boron	5
12.0	<b>C</b>	carbon	6
27.0	<b>Al</b>	aluminium	13
28.1	<b>Si</b>	silicon	14

(14)

12.0	<b>C</b>	carbon	6
14.0	<b>N</b>	nitrogen	7
31.0	<b>P</b>	phosphorus	15
28.1	<b>Si</b>	silicon	14

(15)

14.0	<b>N</b>	nitrogen	7
16.0	<b>O</b>	oxygen	8
32.1	<b>S</b>	sulfur	16
31.0	<b>P</b>	phosphorus	15

(16)

19.0	<b>F</b>	fluorine	9
16.0	<b>O</b>	oxygen	8
35.5	<b>Cl</b>	chlorine	17
32.1	<b>S</b>	sulfur	16

(17)

20.2	<b>Ne</b>	neon	10
19.0	<b>F</b>	fluorine	9
35.5	<b>Cl</b>	chlorine	17
32.1	<b>S</b>	sulfur	16

(18)

4.0	<b>He</b>	helium	2
20.2	<b>Ne</b>	neon	10
39.9	<b>Ar</b>	argon	18

(12)

65.4	<b>Zn</b>	zinc	30
63.5	<b>Cu</b>	copper	29
69.7	<b>Ga</b>	gallium	31
72.6	<b>Ge</b>	germanium	32

(11)

65.4	<b>Zn</b>	zinc	30
63.5	<b>Cu</b>	copper	29
114.8	<b>In</b>	indium	49
118.7	<b>Sn</b>	tin	50

(10)

58.7	<b>Ni</b>	nickel	28
58.9	<b>Co</b>	cobalt	27
106.4	<b>Pd</b>	palladium	46
107.9	<b>Ag</b>	silver	47

(9)

55.8	<b>Fe</b>	iron	26
54.9	<b>Mn</b>	manganese	25
101.1	<b>Ru</b>	ruthenium	44
102.9	<b>Rh</b>	rhodium	45

(8)

55.8	<b>Fe</b>	iron	26
54.9	<b>Mn</b>	manganese	25
101.1	<b>Ru</b>	ruthenium	44
102.9	<b>Rh</b>	rhodium	45

(7)

54.9	<b>Mn</b>	manganese	25
[98]	<b>Tc</b>	technetium	43
106.4	<b>Pd</b>	palladium	46
107.9	<b>Ag</b>	silver	47

(6)

52.0	<b>Cr</b>	chromium	24
95.9	<b>Mo</b>	molybdenum	42
183.8	<b>W</b>	tungsten	74
186.2	<b>Re</b>	rhenium	75

(5)

50.9	<b>V</b>	vanadium	23
92.9	<b>Nb</b>	niobium	41
180.9	<b>Ta</b>	tantalum	73
186.2	<b>Re</b>	rhenium	75

(4)

47.9	<b>Ti</b>	titanium	22
91.2	<b>Zr</b>	zirconium	40
178.5	<b>Hf</b>	hafnium	72
186.2	<b>Re</b>	rhenium	75

(3)

45.0	<b>Sc</b>	scandium	21
88.9	<b>Y</b>	yttrium	39
138.9	<b>La*</b>	lanthanum	57
178.5	<b>Hf</b>	hafnium	72

(2)

137.3	<b>Ba</b>	barium	56
138.9	<b>La*</b>	lanthanum	57
227	<b>Ac*</b>	actinium	89
178.5	<b>Hf</b>	hafnium	72

(1)

6.9	<b>Li</b>	lithium	3
9.0	<b>Be</b>	beryllium	4
23.0	<b>Na</b>	sodium	11
24.3	<b>Mg</b>	magnesium	12

(11)

112.4	<b>Cd</b>	cadmium	48
107.9	<b>Ag</b>	silver	47
114.8	<b>In</b>	indium	49
118.7	<b>Sn</b>	tin	50

(10)

195.1	<b>Pt</b>	platinum	78
197.0	<b>Au</b>	gold	79
200.6	<b>Hg</b>	mercury	80
204.4	<b>Tl</b>	thallium	81

(9)

192.2	<b>Ir</b>	iridium	77
192.2	<b>Os</b>	osmium	76
197.0	<b>Au</b>	gold	79
200.6	<b>Hg</b>	mercury	80

(8)

190.2	<b>Os</b>	osmium	76
192.2	<b>Rh</b>	rhodium	45
195.1	<b>Pt</b>	platinum	78
197.0	<b>Au</b>	gold	79

(7)

186.2	<b>Re</b>	rhenium	75
186.2	<b>Os</b>	osmium	76
192.2	<b>Rh</b>	rhodium	45
195.1	<b>Pt</b>	platinum	78

(6)

186.2	<b>Re</b>	rhenium	75
186.2	<b>Os</b>	osmium	76
192.2	<b>Rh</b>	rhodium	45
195.1	<b>Pt</b>	platinum	78

(5)

180.9	<b>Ta</b>	tantalum	73
180.9	<b>Nb</b>	niobium	41
186.2	<b>Re</b>	rhenium	75
192.2	<b>Ir</b>	iridium	77

(4)

178.5	<b>Hf</b>	hafnium	72
178.5	<b>Zr</b>	zirconium	40
186.2	<b>Re</b>	rhenium	75
192.2	<b>Ir</b>	iridium	77

(3)

178.5	<b>Hf</b>	hafnium	72
178.5	<b>Zr</b>	zirconium	40
186.2	<b>Re</b>	rhenium	75
192.2	<b>Ir</b>	iridium	77

(2)

137.3	<b>Ba</b>	barium	56
138.9	<b>La*</b>	lanthanum	57
227	<b>Ac*</b>	actinium	89
178.5	<b>Hf</b>	hafnium	72

(1)


6.9	<b>Li</b>	lithium	3
9.0	<b>Be</b>	beryllium	4
23.0	<b>Na</b>	sodium	11
24.3	<b>Mg</b>	magnesium	12

Elements with atomic numbers 112-116 have been reported but not fully authenticated

140	<b>Ce</b>	cerium	58	141	<b>Pr</b>	praseodymium	59	144	<b>Nd</b>	neodymium	60	150	<b>Sm</b>	samarium	62	157	<b>Gd</b>	gadolinium	64	163	<b>Dy</b>	dysprosium	66	165	<b>Ho</b>	holmium	67	167	<b>Er</b>	erbium	68	169	<b>Tm</b>	thulium	69	173	<b>Yb</b>	ytterbium	70	175	<b>Lu</b>	lutetium	71
232	<b>Th</b>	thorium	90	231	<b>Pa</b>	protactinium	91	238	<b>U</b>	uranium	92	237	<b>Np</b>	neptunium	93	247	<b>Cm</b>	curium	96	251	<b>Cf</b>	californium	98	254	<b>Es</b>	einsteinium	99	253	<b>Fm</b>	fermium	100	256	<b>Md</b>	mendeleevium	101	254	<b>No</b>	nobelium	102	257	<b>Lr</b>	lawrencium	103

\* Lanthanide series

\* Actinide series



Mob: +974 5526711 / 55249797  
E-mail: rashed.saba@gmail.com

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA

DO NOT WRITE IN THIS AREA



P 6 1 6 5 7 A 0 2 8 2 8